

**CYTOGENETICS IN THREE SPECIES OF
POLYBETES SIMON 1897 FROM ARGENTINA
(ARANEAE, SPARASSIDAE)**

I. KARYOTYPE AND CHROMOSOME BANDING PATTERN

Sergio Gustavo Rodríguez-Gil: Laboratorio de Citogenética y Evolución, Departamento de Ecología, Genética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. Intendente Güiraldes y Costanera Norte, C1428EHA. Ciudad Autónoma de Buenos Aires, Argentina. E-mail: rodrigil@bg.fcen.uba.ar

María Susana Merani: Centro de Investigaciones en Reproducción, Facultad de Medicina, Universidad de Buenos Aires, Paraguay 2155, C1121ABG, Ciudad Autónoma de Buenos Aires, Argentina.

Cristina Luisa Scioscia: División Aracnología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Av. Ángel Gallardo 470, C1405DJR, Ciudad Autónoma de Buenos Aires, Argentina.

Liliana María Mola: Laboratorio de Citogenética y Evolución, Departamento de Ecología, Genética y Evolución, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. Intendente Güiraldes y Costanera Norte, C1428EHA. Ciudad Autónoma de Buenos Aires, Argentina.

ABSTRACT. Species of *Polybetes* are known exclusively from South America. Currently there are 13 described species, 9 occurring in Argentina. Cytogenetic studies in spiders are scarce; the cytogenetics of only about 1% of nearly 39,500 described species are known. Within the Sparassidae, 38 species out of 1,009 (< 4%) have been cytogenetically analyzed; the most frequent chromosome number is $2n = 43/46$ (male/female), $n = 20 + X_1X_2X_3$, present in almost half of the species studied. Female diploid chromosome number is only known for four species: *Heteropoda venatoria* (Linnaeus 1767) ($2n = 44$); *Pediana regina* (L. Koch 1875), *Isopeda* sp. and *Olios* sp. ($2n = 46$). Within the genus *Polybetes*, only *P. pythagoricus* (Holmberg 1875) had been previously cytogenetically analyzed. In the present work, the karyotype, heterochromatin content and distribution, and silver stained nucleolus-organizer regions of *P. pythagoricus*, *P. rapidus* (Keyserling 1880) and *P. punctulatus* Mello-Leitão 1944 are described and compared. In *P. pythagoricus* the identification of the chromosome pairs by means of G-banding is also performed. Females of the three species show a chromosome complement of 44 telocentric chromosomes, with a similar karyotype. Males of *P. pythagoricus* show 42 telocentric chromosomes, the two sex chromosomes being the largest and of different size. In the three species, two pairs of telomeric NORs and small pericentromeric positive C-bands in all chromosomes were detected. This C-banding pattern seems to be characteristic of spiders. Comparative analysis of chromosome complements in Sparassidae indicates that $2n = 42/44$ ($X_1X_20/X_1X_1X_2X_2$) (male/female) may represent the ancestral karyotype for *Polybetes*.

Keywords: Chromosome number, telocentric chromosomes, heterochromatin, nucleolus-organizing regions

Species of *Polybetes* are known exclusively from South America. To date there are thirteen described species, nine of them occurring in Argentina (Platnick 2006): *P. germaini* Simon 1897, *P. martius* (Nicolet 1849), *P. obnuptus* Simon 1896, *P. pallidus* Mello-Leitão 1941, *P. punctulatus* Mello-Leitão 1944, *P. pythagoricus* (Holmberg 1875), *P. quadrifoveatus*

(Järvi 1914), *P. rapidus* (Keyserling 1880), and *P. trifoveatus* (Järvi 1914). In nature, they are found under the bark of trees (e.g., *P. pythagoricus* is common under the bark of *Eucalyptus*), in the branches of trees (*P. rapidus*), and others are found in grasses such as *Cortaderia* spp. (*P. punctulatus*). *Polybetes pythagoricus* and *P. rapidus* are also common in

cities where they are found in parks, gardens or even in the roofs of buildings. They are nocturnal and sometimes enter houses on stormy days. Despite their usual aggressiveness, they possess venom of low toxicity that causes little local injury and is harmless to humans (Scioscia, personal observations).

Cytogenetic studies in spiders are scarce, with only approximately 1% of nearly 39,500 described species determined. Within the Sparassidae, 36 species out of 1,009 (< 4%) had been previously cytogenetically analyzed; male diploid chromosome numbers range from 21 to 44; female diploid chromosome number is only known for four species: *Heteropoda venatoria* (Linnaeus 1767) ($2n = 44$); *Pediana regina* (L. Koch 1875), *Isopeda* sp. and *Olios* sp. ($2n = 46$). The most frequent sex chromosome determination system is $X_1X_2X_3O/X_1X_1X_2X_3X_3$ (male/female) (Hackman 1948; Suzuki 1950; Suzuki & Okada 1950; Bole-Gowda 1952; Suzuki 1952; Mittal 1961; Díaz & Sáez 1966a, b; Mittal 1966; Benavente & Wettstein 1978; Olivera 1978; Datta & Chatterjee 1983; Rowell 1985; Srivastava & Shukla 1986; Parida & Sharma 1986, 1987; Rowell 1991a, b; Hancock & Rowell 1995; Platnick 2006). Within the genus *Polybetes*, only *P. pythagoricus* had been previously cytogenetically analyzed (Díaz & Sáez 1966a, b; Benavente & Wettstein 1978; Olivera 1978).

There are only a few studies in spiders that characterize banding patterns of chromosomes; the distribution of C heterochromatin is known in eleven Sparassidae, eight Araneidae, five Lycosidae, four Tetragnathidae, two Nephilidae, two Sicariidae, one Scytodidae, and one Salticidae species, and G-banding was performed only in *Lycosa thorelli* (Keyserling 1877) (Lycosidae) and *P. pythagoricus* (Brum-Zorrilla & Cazenave 1974; Olivera 1978; Brum-Zorrilla & Postiglioni 1980; Rowell 1985; Datta & Chatterjee 1988; Rowell 1991b; Gorlova et al. 1997; Silva et al. 2002; De Araujo et al. 2005a, b).

In the present work, the karyotype, heterochromatin content and distribution, and silver stained nucleolus-organizer regions (NORs) of *Polybetes pythagoricus*, *P. rapidus* and *P. punctulatus* are described and compared. Furthermore, in *P. pythagoricus* the identification of the chromosome pairs by means of G-banding was performed.

METHODS


Adult females and males were collected in the field and were bred at the Arachnology Division of the Museo Argentino de Ciencias Naturales "Bernardino Rivadavia" (MACN). Voucher specimens of all species in this study have been deposited in the National Collection of Arachnology (MACN-Ar, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Cristina Scioscia): *Polybetes pythagoricus*, five females and three males from Buenos Aires City, 34°35'15"S, 58°40'21"W, and Buenos Aires Province (Los Polvorines, 34°30'00"S, 58°41'00"W; Villa Madero, 34°42'00"S, 58°30'00"W; San Isidro, 34°28'15"S, 58°31'43"W; and Martín García Island Natural Preserve, 34°11'15"S, 58°16'52"W); *P. rapidus*, six females from Buenos Aires Province (Bella Vista, 35°14'00"S, 59°53'00"W; Merlo, 34°40'12"S, 58°45'10"W; Villa Madero and Martín García Island Natural Preserve); *P. punctulatus*, one female from Martín García Island Natural Preserve and two immature females born in the lab.

For cytogenetic preparations, the specimens were cooled; and injected with 0.1 ml of 0.01% colchicine solution. After 1.25 h, several drops of hemolymph were removed from the coxal joints, and gonads together with some digestive tissues were dissected. Each sample was dispersed in 2 ml of hypotonic solution (KCl 0.56%) for 15 min, centrifuged at 800 rpm for 5 min, and fixed in 1 ml of 3:1 (methanol:acetic acid). The cell suspension was dropped onto clean slides, air-dried and stained with Giemsa 3% for chromosome counts and karyotyping.

C-band preparations were made following Sumner (1972) with some modifications: 0.2 N HCl for 1 h; saturated solution of Ba(OH)₂ for 30 s–1 min; 2 × SSC at 60° C for 1 h. Slides were air-dried and stained with 3% Giemsa.

G-bands preparations were made as follows: PBS solution for 15 min; 0.1% trypsin for 45 s–1 min. Slides were air-dried and stained with 3% Giemsa. NOR-banding was performed following Howell & Black (1980).

Eight to fifteen well-spread mitotic metaphases were measured to determine the karyotype of each species. Chromosome measurements were made using the computer



Figures 1–4.—Mitotic metaphases in spider chromosomes. 1. *Polybetes rapidus* female ($2n = 44$); 2. *P. punctulatus* female ($2n = 44$); 3. *P. pythagoricus* female ($2n = 44$); 4. *P. pythagoricus* male ($2n = 42$). Scale = 10 μm .

application Micromesure version 3.3 (Reeves & Tear 2000). The total haploid complement length (TCL) in females was calculated by adding the mean value of each chromosome pair (in arbitrary units). In males of *P. pythagoricus*, the relative length of all chromosomes was analyzed to identify the two chromosomes that have no homologues (sex chromosomes), and TCL was afterwards calculated. The idiogram of each species was drawn on the basis of the relative percentage of each chromosome pair length to the TCL. Chromosome measurements were also made using a vernier calliper to estimate TCL in microns.



RESULTS

Chromosome complement.—Females of *P. rapidus*, *P. punctulatus* and *P. pythagoricus* show a chromosome complement of 44 telocentric chromosomes, and 42 telocentric

chromosomes in males of *P. pythagoricus*. The sex chromosomes cannot be distinguished by their differential pycnosis in somatic metaphases of males and females (Figs. 1–4).

The total haploid complement length (TCL) is similar in the three species: $67.29 \pm 4.91 \mu\text{m}$ in *P. rapidus*, $66.28 \pm 6.91 \mu\text{m}$ in *P. pythagoricus* and $63.70 \pm 1.52 \mu\text{m}$ in *P. punctulatus*.

Females of the three species show a similar karyotype: there are three large pairs of differently-sized chromosomes that can be distinguished; the rest of the chromosomal complement gradually decreases in size, except for the last pair that is slightly smaller. The largest chromosome pair shows slight size differences in the three species (*P. pythagoricus*, 7.00%; *P. punctulatus*, 6.80%; and *P. rapidus*, 6.63%), while the second pair is longer in *P. pythagoricus* (6.45%) (*P. punctulatus*, 6.12%; and *P. rapidus*, 6.16%) (Figs. 5, 7–9). In




Figures 5, 6.—Comparative ideograms of spider chromosomes. 5. Females of *Polybetes rapidus*, *P. punctulatus* and *P. pythagoricus*; 6. Female and male of *P. pythagoricus*.

males of *P. pythagoricus*, the length analysis of all chromosomes of the complement makes it possible to determine that the two largest chromosomes of different sizes are the sex chromosomes X_1X_2 (Figs. 6, 10). Meiotic studies performed in males of *P. punctulatus* and *P. rapidus* (Rodríguez Gil 2006) demonstrated that in these species the sex chromosomes are also the largest of the complement.

C-banding and NORs silver staining.—In females of the three species, small positive

C-bands in the pericentromeric region of all chromosomes were detected, except in the X_2 pair of *P. pythagoricus* where they are more prominent (Figs. 11, 12). The number of chromosomes with telomeric nucleolus-organizer regions (NORs) silver stained varied from 1 to 4 in different cells of the three species (Figs. 13, 14). It was not possible to determine precisely the NOR pairs; one pair was medium sized and the other was among the smaller ones.



Figures 7–10.—Karyograms from spider cells depicted in Figures 1–4. 7. *Polybetes rapidus* female; 8. *P. punctulatus* female; 9. *P. pythagoricus* female; 10. *P. pythagoricus* male.


G-banding.—Despite applying the G-banding technique to the three species, only *P. pythagoricus* females yielded reproducible results, making it possible to determine the chromosome pair's identification (Figs. 16, 17). In *P. punctulatus*, only a few dark bands, with little contrast, were obtained in all the chromosomes (Fig. 15); therefore, the chromosome pair's identification was not possible. In *P. rapidus* no bands were obtained.

DISCUSSION

Chromosome complement and karyotype.—*Polybetes punctulatus* and *P. rapidus* were here cytogenetically characterized for the first time. *Polybetes pythagoricus* had been previously analyzed in Uruguayan populations; Benavente & Wettstein (1978) performed an ultrastructural study of sex chromosomes pairing in meiosis, and Díaz & Sáez (1966a, b) reported $2n = 42$ and $n = 20 + X_1X_2$ in males. However, Olivera (1978), in a preliminary report of *P. pythagoricus*, described contradictory data reporting a $2n = 40/42$ (male/female) with sex determination system $X_1X_2/X_1X_1X_2X_2$ at mitosis but in male meiosis described the presence of 10 chromosomes plus 2 Xs at one pole and 10 chromosomes at the other one in anaphase I. The three species of *Polybetes* analyzed in this work have $2n = 44 = 40 + X_1X_1X_2X_2$ in females and $2n = 42 = 40 + X_1X_2$ in *P. pythagoricus* males; they show very similar karyotype and total haploid complement length,

with all the chromosomes telocentric. The sex chromosomes are the largest ones, the X_1 show slight size differences in the three species and X_2 is longer in *P. pythagoricus*. The conservative karyotype present in the three species could be considered characteristic for the genus.

Currently, cytogenetic studies in Sparassidae have been performed on 38 species from 17 genera (Table 1). Usually the chromosomes are telocentric and one of the sex chromosomes is the largest of the complement. The predominant diploid numbers in this family are $2n = 43$, $n = 20 + X_1X_2X_3$, 15 species; and $2n = 41$, $n = 19 + X_1X_2X_3$, 10 species. In other genera, besides *Polybetes*, there seems to be karyotypic conservation as in *Heteropoda* ($n = 19 + X_1X_2X_3$, in 5 of 6 species studied) and *Isopeda* ($n = 20 + X_1X_2X_3$, in the 4 species analyzed). Bole-Gowda (1952) stated that *Heteropoda sexpunctata* Simon 1885 has a derived karyotype, $2n = 20 + X$ (male) with 19 metacentric (including the X chromosome) and two acrocentric autosomes, on the basis of Robertsons law. On the other hand, in the genus *Sparassus*, there is variation not only in chromosome numbers ($2n = 22$ to 44) but in the sex chromosome determination system as well (X_1X_2 , $X_1X_2X_3$, $X_1X_2X_3X_4$); although none of the entities was identified at the species level, and it is possible that the genus may be misidentified in some of them (Parida & Sharma 1987). An-



Figures 11-16.—C-banding in spider chromosomes: 11. *Polybetes punctulatus* female; 12. *P. pythagoricus* female (arrowheads point to prominent C-bands). NORs silver staining in spider chromosomes: 13. *P. rapidus* female; 14. *P. pythagoricus* female (arrows point to NOR regions). G-banding in spider chromosomes: 15. *P. punctulatus* female; 16. *P. pythagoricus* female. Scale = 10 μ m.




Figure 17.—G-banding karyogram and idiogram of *Polybetes pythagoricus* chromosomes (from cell depicted in Fig. 16).

cestral populations of *Delena cancerides* Walckenaer 1837 also show $n = 20 + X_1X_2X_3$, but this species has a number of chromosomal races that differ by the presence of particular combinations of chromosomal fusions, either in homozygous or heterozygous condition (Rowell 1985, 1990, 1991a, b; Hancock & Rowell 1995). A reduced complement is also present in *Micrommata virescens* (Clerck 1757), but neither the chromosome number nor the sex chromosome complement is known with certainty (Hackman 1948).

Heterochromatin characterization.—The three species of *Polybetes* analyzed here show only small pericentromeric heterochromatic bands in all the chromosomes with no differential pycnosis of the sex chromosomes, although Olivera (1978) reported that “substantial” heterochromatic blocks were present in *P. pythagoricus* mitotic and meiotic chromosomes. *Polybetes pythagoricus* X_2 chromosomes showed a larger C-band than in the other two species, which may explain the differences in these chromosomes’ size.

Since the pioneer characterization of *Schizocosa malitiosa* (Tullgren 1905) (Lycosidae)

by Brum-Zorrilla & Cazenave (1974), few spider species have been analyzed with regard to the heterochromatin content and distribution. Our results fit with previous data of most of the other spider species analyzed that have a small amount of pericentromeric heterochromatin in the autosomes and sex chromosomes; this condition seems to be characteristic in spiders. In *Loxosceles intermedia* Mello-Leitão 1934 (Sicariidae) and *Isopeda* species, pericentromeric C-bands are more conspicuous (Brum-Zorrilla & Postiglioni 1980; Rowell 1985, 1991b; Datta & Chatterjee 1988; Gorlova et al. 1997; Silva et al. 2002). In a few species, telomeric localization of heterochromatin (telomeric C-bands) has also been described in some chromosomes of the complement; these bands have usually appeared in a polymorphic condition (Rowell 1985; Datta & Chatterjee 1988; Rowell 1991b; De Araujo 2005a). In *Nephilengys cruentata* (Fabricius 1775) (Nephilidae) interstitial C-bands are present in some autosomes; the same occurs in some autosomes and the sex chromosomes of one unidentified species of *Scytodes* (De Araujo et al. 2005a, b).

Table 1.—Karyotype characteristics and collecting locality of the Sparassidae species cytogenetically analyzed (f = female).

Species	2n	n (male)	Locality	References
<i>Bhutaniella sikkimensis</i> (Gravely 1931)	42	19 + X ₁ X ₂ X ₃ X ₄	India	Datta & Chatterjee 1983
<i>Delena cancerides</i> Walckenaer 1837 (ancestral karyotype)	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1985, 1991a, b; Hancock & Rowell 1995
<i>Delena</i> sp.	43	20 + X ₁ X ₂ X ₃		McIntosh in Suzuki 1952
<i>Heteropoda leprosa</i> Simon 1884	41	19 + X ₁ X ₂ X ₃	India	Datta & Chatterjee 1983
<i>Heteropoda phasma</i> Simon 1897	41	19 + X ₁ X ₂ X ₃	India	Srivastava & Shukla 1986
<i>Heteropoda procera</i> (L. Koch 1867)	41	19 + X ₁ X ₂ X ₃	Australia	Rowell 1985
<i>Heteropoda sexpunctata</i> Simon 1885	21	10 + X	India	Bole-Gowda 1952
<i>Heteropoda venatoria</i> (Linnaeus 1767)	41–44 f	19 + X ₁ X ₂ X ₃	India, Japan	Suzuki & Okada 1950; Bole-Gowda 1952; Srivastava & Shukla 1986
<i>Heteropoda</i> sp. nov.	41	19 + X ₁ X ₂ X ₃	Australia	Rowell 1985
<i>Holconia immanis</i> (L. Koch 1867)	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991a, b (sub <i>Isopoda</i>)
<i>Isopoda vasta</i> (L. Koch 1867)	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991b (sub <i>Isopoda vaster</i> (sic))
<i>Isopoda villosa</i> L. Koch 1875	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991a, b (sub <i>Isopoda</i>)
<i>Isopoda</i> sp.	43–46 f	20 + X ₁ X ₂ X ₃	Australia	Rowell 1985 (sub <i>Isopoda</i>)
<i>Isopoda</i> sp. nov.	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991b (sub <i>Isopoda</i>)
<i>Isopedella leai</i> Hogg 1903	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991b (sub <i>Isopoda tepperi</i> Hogg)
<i>Micrommata virescens</i> (Clerck 1757)	±35	16 + X ₁ X ₂ X ₃ (?)	Finland	Hackman 1948 [sub <i>Micrommata viridissima</i> (De Geer)]
<i>Neosparassus diana</i> (L. Koch 1875)	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991b (sub <i>Olios</i>)
<i>Olios lamarcki</i> (Latreille 1806)	42	20 + X ₁ X ₂	India	Bole-Gowda 1952
<i>Olios</i> sp. 1	43	20 + X ₁ X ₂ X ₃		McIntosh in Suzuki 1952
<i>Olios</i> sp. 2	43–46 f	20 + X ₁ X ₂ X ₃	Australia	Rowell 1985
<i>Parapalystes whiteae</i> (Pocock 1902)	43	20 + X ₁ X ₂ X ₃	India	Mittal 1961, 1966 (sub <i>Palystes</i>)
<i>Pediana regina</i> (L. Koch 1875)	43–46 f	20 + X ₁ X ₂ X ₃	Australia	Rowell 1985, 1991b
<i>Pediana</i> sp. nov.	43	20 + X ₁ X ₂ X ₃	Australia	Rowell 1991b
<i>Polybetes punctulatus</i> Mello-Leitão 1944	44 f	20 + X ₁ X ₂	Argentina	this work; Rodríguez Gil 2006
<i>Polybetes pythagoricus</i> (Holmberg 1875)	42–44 f	20 + X ₁ X ₂	Uruguay	Díaz & Sáez 1966a, b (sub <i>P. pitagorica</i> (sic))
			Argentina	this work; Rodríguez Gil 2006
	40–42 f		Uruguay	Olivera 1978 (sub <i>P. pythagoricus</i> (sic))
<i>Polybetes rapidus</i> (Keyserling 1880)	44 f	20 + X ₁ X ₂	Argentina	this work; Rodríguez Gil 2006

Table 1.—Continued.

Species	2n	n (male)	Locality	References
<i>Pseudopoda prompta</i> (O. P.-Cambridge 1885)	41	19 + X ₁ X ₂ X ₃	India	Srivastava & Shukla 1986 (sub <i>Heteropoda</i>)
<i>Sinopoda forcipata</i> (Karsch 1881)	41	19 + X ₁ X ₂ X ₃	Japan	Suzuki 1952 (sub <i>Heteropoda</i>)
<i>Sparassus</i> sp. 1	44	21 + X ₁ X ₂	India	Parida & Sharma 1987
<i>Sparassus</i> sp. 2	42	20 + X ₁ X ₂	India	Parida & Sharma 1987
<i>Sparassus</i> sp. 3	41	19 + X ₁ X ₂ X ₃	India	Parida & Sharma 1986, 1987
<i>Sparassus</i> sp. 4	41	19 + X ₁ X ₂ X ₃	India	Parida & Sharma 1987
<i>Sparassus</i> sp. 5	22	10 + X ₁ X ₂	India	Parida & Sharma 1987
<i>Sparassus</i> sp. 6	42	20 + X ₁ X ₂	India	Datta & Chatterjee 1983 (sub <i>Parassus</i> sp. 1)
<i>Sparassus</i> sp. 7	44	20 + X ₁ X ₂ X ₃ X ₄	India	Datta & Chatterjee 1983 (sub <i>Parassus</i> sp. 2)
<i>Sparassus</i> sp. 8	42	20 + X ₁ X ₂	India	Datta & Chatterjee 1983
<i>Spariolenus tigris</i> Simon 1880	41	19 + X ₁ X ₂ X ₃	India	Bole-Gowda 1952
<i>Thelcticopis severa</i> (L. Koch 1875)	43	Possibly X ₁ X ₂ X ₃	Japan	Suzuki 1950, 1952 (sub <i>Thelcticopis</i> (sic))

In spermatogonial mitosis, after C-banding, sex chromosomes have shown two different patterns. In three species of Lycosidae and in *Delena cancerides* the sex chromosomes were more darkly stained than the autosomes, while there was no difference in the sex chromosomes and autosomes in *Isopeda* and Araneidae species. In the three *Polybetes* species presented here and in four araneids, there is also no difference in the sex chromosomes and autosomes in female somatic and gonial cells. In one species, *Schizocosa malitiosa*, only one X chromosome was notable in that it exhibited complete heterochromatinization (Brum-Zorrilla & Cazenave 1974; Brum-Zorrilla & Postiglioni 1980; Rowell 1985; Datta & Chatterjee 1988; Rowell 1991b).

NORs silver staining and G-banding.—The variation in the number of chromosomes per cell bearing nucleolus-organizer regions observed in *Polybetes* species is common in the Ag-technique. It is characteristic of silver staining that not all the NORs are usually silver stained in species with multiple NORs, but only those transcriptionally active during the preceding interphase (Sumner 2003). It can be concluded that two chromosomal pairs with telomeric NORs are present in the three species here analyzed. Although the identification of the NOR pairs should be regarded as tentative, it seems possible that they correspond

to the same pairs in the three species. Only a pair of NORs was detected at early somatic stages of *P. pythagoricus* Uruguayan specimens (Olivera 1978).

G-banding allows the precise identification of homologues and facilitates karyotypic comparisons between related species. Although good quality G-bands can be produced in reptiles, birds, mammals, in some fishes and amphibians, and in a few plants, this method does not yield consistent results in invertebrate chromosomes and only in a few species of insects have well-defined G-bands been obtained. The difficulty in obtaining good quality G-bands in invertebrates may reflect differences in mitotic chromosome substructure, e.g., tight compaction of the chromatin compared to vertebrates (Lorite et al. 1996; Appels et al. 1998; Baldanza et al. 1999; Sumner 2003). In spiders, G-banding had been performed in three species of Lycosidae (but only *Lycosa thorelli* showed consistent G-banding), and in *P. pythagoricus* from Uruguay (where only a few pairs could be identified) (Olivera 1978; Brum-Zorrilla & Postiglioni 1980). In the present work, the identification of all chromosome pairs was possible in *P. pythagoricus*. Taking into account that the pattern of pachytene chromomeres resembles that of G-bands on the same chromosome (Sumner 2003), it would be interesting to perform a

comparative analysis of the chromomere pattern of the three species in order to know if the scarcity and absence of G-bands in *P. punctulatus* and *P. rapidus* respectively is due to structural differences or to technical procedures.

The three species of *Polybetes* here analyzed are easily distinguished by morphological characters, but they are very conservative karyotypically. This fact could be useful in future for the delimitation of genera in a systematic revision of the family.

ACKNOWLEDGMENTS

The present study was supported by grants from the National University of Buenos Aires (UBA) (Ex 317 to Drs. L. Poggio and L. Mola) and from the National Council of Scientific and Technological Research (CONICET) (PIP 02296 and 05927 to Drs. L. Poggio and L. Mola and PIP 02202 and 05654 to Drs. A. González and C. Scioscia). The authors wish to thank to Mr. Hernán Dinapoli for technical assistance, to Lic. Pablo Rebagliati, Lic. Mariana López and the student Luis Piacentini for collecting some of the specimens, and to Dr. María Inés Pigozzi for critical reading of the manuscript. The authors wish to dedicate this paper to the memory of Dr. Carlos A. Naranjo.

LITERATURE CITED

- Appels, R., R. Morris, B.S. Gill & C.E. May. 1998. Chromosome Biology. Kluwer Academic Publishers, Boston, Massachusetts. 401 pp.
- Baldanza, F., L. Gaudio & G. Viggiani. 1999. Cytotaxonomic studies of *Encarsia* Förster (Hymenoptera: Aphelinidae). Bulletin of Entomological Research 89:209–215.
- Benavente, R. & R. Wettstein. 1978. Ultrastructural cytogenetics of the sex determination mechanisms of araneids. Revista de Microscopía Electrónica 5:320–321.
- Bole-Gowda, B.N. 1952. Studies on the chromosomes and the sex-determining system in four hunting spiders (Sparassidae). Proceedings of the Zoological Society, Bengal 5:51–70.
- Brum-Zorrilla, N. & A. Postiglioni. 1980. Kariological studies on Uruguayan spiders. I. Banding pattern in chromosomes of *Lycosa* species (Araneae-Lycosidae). Genetica 54:149–153.
- Brum-Zorrilla, N. & A.M. Cazenave. 1974. Heterochromatin localization in the chromosomes of *Lycosa malitiosa* (Arachnida). Experientia 30: 94–95.
- Datta, S.N. & K. Chatterjee. 1983. Chromosome number and sex-determining system in fifty-two species of spiders for north-east India. Chromosome Information Service 35:6–8.
- Datta, S.N. & K. Chatterjee. 1988. Chromosome and sex determination in 13 araneid spiders of North-Eastern India. Genetica 76:91–99.
- De Araujo, D., D.M. Cella & A.D. Brescovit. 2005a. Cytogenetic analysis of the neotropical spider *Nephilengys cruentata* (Araneomorphae, Tetragnathidae): Standard staining, Nors, C-Bands and base-specific fluorochromes. Brazilian Journal of Biology 65:193–202.
- De Araujo, D., D.M. Cella & A.D. Brescovit. 2005b. Variabilidade cromossômica em três espécies de *Scytodes* Latreille, 1804 (Araneae, Haplogynae, Scytodidae). Primer Congreso Latinoamericano de Aracnología (Minas, Uruguay) Actas: 107.
- Díaz, M. & F.A. Sáez. 1966a. Karyotypes of South American Araneida. Memórias do Instituto Butantan Commemorativo 33:153–154.
- Díaz, M. & F.A. Sáez. 1966b. Investigaciones citogenéticas sobre algunas especies de araneidos uruguayos. Anales (II) Congreso Latinoamericano de Zoología (San Pablo):3–9.
- Gorlova, O.Y., J.P. Gorlov, E. Nevo & D.V. Logunov. 1997. Cytogenetic studies on seventeen spider species from Israel. Bulletin of the British Arachnological Society 10:249–252.
- Hackman, W. 1948. Chromosomenstudien an Araneen mit besonderer Berücksichtigung der Geschlechtschromosomen. Acta Zoologica Fennica 54: 1–101.
- Hancock, A.J. & D.M. Rowell. 1995. A chromosomal hybrid zone in the Australian huntsman spider, *Delena cancerides* (Araneae: Sparassidae). Evidence for a hybrid zone near Canberra, Australia. Australian Journal of Zoology 43:173–180.
- Howell, W.M. & D.A. Black. 1980. Controlled silver staining of nucleolus organizer regions with a protective colloidal developer: a 1-step method. Experientia 36:1014–1015.
- Lorite, P., E. Chica & T. Palomeque. 1996. G-banding and chromosome condensation in the ant, *Tapinoma nigerrimum*. Chromosome Research 4:77–79.
- Mittal, O.P. 1961. Chromosome number and sex mechanism in twenty-one species of the Indian spiders. Research Bulletin (N. S.) of the Panjab University 12:271–273.
- Mittal, O.P. 1966. Karyological studies on Indian spiders. IV. Chromosomes in relation to taxonomy in Eusparassidae, Selenopidae and Thomisidae. Genetica 37:225–234.
- Olivera, G. 1978. Estudio citogenético en *Polybetes pythagoricus* (Araneomorphae-Sparassidae). XX. Semana Univeritaria Gaucha de Debates Biológicos, Porto Alegre 24(Abstracts):251–253.
- Parida, B.B. & G.P. Sharma. 1986. Karyotype and

- spermatogonia in an Indian hunting spider, *Sparassus* sp. (Sparassidae: Arachnida) with multiple sex chromosomes. *Chromosome Information Service* 40:28–30.
- Parida, B.B. & G.P. Sharma. 1987. Chromosome number, sex mechanism and genome size in twenty seven species of Indian spiders. *Chromosome Information Service* 43:11–13.
- Platnick, N.I. 2006. The World Spider Catalog, version 7.0. American Museum of Natural History, New York. Online at <http://research.amnh.org/entomology/spiders/catalog/index.html>.
- Reeves, A. & J. Tear. 2000. Micromasure for Windows. Version 3.3 free program distributed by the author from <http://www.colostate.edu/Depts/Biology/Micromasure>.
- Rodríguez-Gil, S.G. 2006. Citogenética de arañas sinantrópicas de Argentina (Arachnida: Araneae). Ph. D. Thesis. Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires. 155 pp.
- Rowell, D.M. 1985. Complex sex-linked fusion heterozygosity in the Australian huntsman spider *Delena cancerides* (Araneae: Sparassidae). *Chromosoma* (Berlin) 93:169–176.
- Rowell, D.M. 1990. Fixed fusion heterozygosity in *Delena cancerides* Walck. (Araneae: Sparassidae): an alternative to speciation by monobranched fusion. *Genetica* 80:139–157.
- Rowell, D.M. 1991a. Chromosomal fusion and meiotic behaviour in *Delena cancerides* (Araneae: Sparassidae). I. Pairing behaviour and X-chromosome segregation. *Genome* 34:561–566.
- Rowell, D.M. 1991b. Chromosomal fusion and meiotic behaviour in *Delena cancerides* (Araneae: Sparassidae). II. Chiasma position and its implications for speciation. *Genome* 34:567–573.
- Silva, R.W., D.D.R. Klisiowicz, D.M. Cella, O.C. Mangili & I.J. Sbalqueiro. 2002. Differential distribution of constitutive heterochromatin in two species of brown spider: *Loxosceles intermedia* and *L. laeta* (Araneae, Sicariidae), from the metropolitan region of Curitiba, PR (Brazil). *Acta Biológica Paranaense, Curitiba* 31:123–136.
- Srivastava, S.C. & S. Shukla. 1986. Chromosome number and sex determining mechanism in forty-seven species of Indian spiders. *Chromosome Information Service* 41:23–26.
- Sumner, A.T. 1972. A simple technique for demonstrating centromeric heterochromatin. *Experimental Cell Research* 75:304–305.
- Sumner, A.T. 2003. *Chromosomes: Organization and Function*. Blackwell Publishing, Oxford, UK. 287 pp.
- Suzuki, S. 1950. Sex determining mechanism and karyotypes in spiders. *Zoological Magazine* (Tokyo) 59:31–32.
- Suzuki, S. 1952. Cytological Studies in spiders II. Chromosomal investigation in the twenty two species of spiders belonging to the four families, Clubionidae, Sparassidae, Thomisidae, and Oxyopidae, which constitute Clubionoidea, with special reference to sex chromosomes. *Journal of Science of the Hiroshima University, Series B, Division. 1* 13:1–52.
- Suzuki, S. & A. Okada. 1950. A study on the chromosomes of a spider, *Heteropoda venatoria*, with special reference to X1-, X2- and X3- chromosomes. *Journal of Science of the Hiroshima University, Series B, Division 1*. 11:1–47.

Manuscript received 2 August 2005, revised 19 January 2007.